Author Index

ABATANGELO, C., PLOTKIN, L., MATHOV, I., SQUIQUERA, L. & LEONI, J. The frequent mutation Gly/Asp in CDR1H may determine a cross-reactive idiotope in anti-I cold agglutinins, 185
ABE, A. *see* TAKAHASHI, T.
ADAMS, D.H. *see* COCKWELL, P.
ADOLF, M. *see* KUCHARZIK, T.
AHLFORS, E., JONSSON, R. & CZERKINSKY, C. Experimental T cell-mediated inflammatory reactions in the murine oral mucosa.II.Immunohistochemical characterization of resident and infiltrating cells, 297
AHO, K. *see* VAARALA, O.
ÅKERBLOM, H.K. *see* LÖNNROT, M.
ALEXANDER, G.J.M. *see* WONG, V.S.
ALFTHAN, G. *see* VAARALA, O.
ALLEN, P.J. *see* HOGARTH, M.B.
ALSINA, J. *see* MESTRE, M.
AMEMIYA, E. *see* NAKAYAMA, T.
ANDERSON, A.C. *see* LI, J.-M.
ANDERSSON, R. *see* ESIN, S.
ANTONELLI, G., GIANNELLI, G., CURRENTI, M., SIMEONI, E., DEL VECCHIO, S., MAGGI, F., PISTELLO, M., ROFFI, L., PASTORE, G., CHEMELLO, L. & DIANZANI, F. Antibodies to interferon (IFN) in hepatitis C patients relapsing while continuing recombinant IFN-α2 therapy, 384
AOYAGI, T. *see* MATSUOKA, N.
AOYAGI, T. *see* MIGITA, K.
APT, A.S. *see* NIKONENKO, B.V.
ARAKAWA, M. *see* SUZUKI, S.
ARANA, R. *see* BACMAN, S.
ARVILOMMI, H. *see* KANTELE, J.M.
ÅSJÖ, B. *see* RØSOK, B.
AUTEGARDEN, J.-E. *see* KEKWICK, R.
AVDIENKO, V.G. *see* NIKONENKO, B.V.
AVRAMEAS, S. *see* MATSIOTA-BERNARD, P.
AXELSSON, M. *see* RØSOK, B.
AGUAS, A.P. *see* ESAGUY, N.

BACMAN, S., STERIN-BORDA, L., JOSÉ CAMUSSO, J., ARANA, R., HUBSCHER, O. & BORDA, E. Circulating antibodies against rat parotid gland M_3 muscarinic receptors in primary Sjögren's syndrome, 454
BADILLO-FLORES, A. *see* MÉNDEZ-SAMPERIO, P.
BANCHEREAU, J. *see* FRIMAN, V.
BANCHEREAU, J. *see* MENETRIER-CAUX, C.
BAROT-CIORBARU, R. *see* TUČKOVÁ, L.
BAS, J. *see* MESTRE, M.
BATONI, G. *see* ESIN, S.
BATSFORD, S.R. *see* MORIOKA, T.
BAYER, B. *see* HOFFMANN, J.C.
BERTRAND, G. *see* RAJAGOPALAN-LEVASSEUR, P.
BEYNON, H.L.C. *see* BOTTO, M.
BHAMBRI, S. *see* KEKWICK, R.
BIRD, C. *see* WADHWA, M.
BJERKNES, R. *see* RØSOK, B.
Borda, E. *see* Bacman, S.
BOTTO, M., THEODORIDIS, E., THOMPSON, E.M., BEYNON, H.L.C., BRIGGS, D., ISENBERG, D.A., WALPORT, M.J. & DAVIES, K.A. FcγRIIa polymorphism in systemic lupus erythematosus (SLE): no association with disease, 264
BRIDON, J.-M. *see* FRIMAN, V.
BRIERE, F. *see* MENETRIER-CAUX, C.
BRIÈRE, F. *see* FRIMAN, V.
BRIGGS, D. *see* BOTTO, M.
BROIDE, D.H. *see* OHNISHI, T.
BROUET, J.C. *see* LEVY, Y.
BROUG-HOLUB, E. & KRAAL, G. Dose- and time-dependent activation of rat alveolar macrophages by glucocorticoids, 332
BROWN, D. *see* WONG, V.S.
BRUIJN, J.A. *see* LUCA, M.E.
BUENDIA, E. *see* MESTRE, M.
BUREK, C.L. *see* JAUME, J.C.

CAMPA, M. *see* ESIN, S.
CAPRON, M. *see* DESREUMAUX, P.
CARROLL, S.F. *see* STOFFEL, M.P.
CARTER, R. *see* WIJESEKERA, S.K.
CASADEMONT, J. *see* CID, M.-C.
CASTELAO, A.M. *see* MESTRE, M.
CAYOTA, A., VUILLIER, F., GONZALEZ, G. & DIGHIERO, G. $CD4^+$ lymphocytes from HIV-infected patients display impaired CD45-associated tyrosine phosphatase activity which is enhanced by anti-oxidants , 11
CHABANE, M.H. *see* KEKWICK, R.
CHAPMAN, K.R. *see* MOLFINO, N.A.
CHARLESWORTH, J.A. *see* GREENSTEIN, J.D.
CHATURVEDI, U.C. *see* MISRA, A.
CHEMELLO, L. *see* ANTONELLI, G.
CHEN, G., WILSON, R., WANG, S.H., ZHENG, H.Z., WALKER, J.J. & MCKILLOP, J.H. Tumour necrosis factor-alpha (TNF-α) gene polymorphism and expression in pre-eclampsia , 154
CHIU, B. *see* INMAN, R.D.
CID, M.-C., GRAU, J.-M., CASADEMONT, J., TOBÍAS, E., PICAZO, A., COLL-VINENT, B., ESPARZA, J., PEDROL, E. & URBANO-MÁRQUEZ, A. Leucocyte/endothelial cell adhesion receptors in muscle biopsies from patients with idiopathic inflammatory myopathies (IIM), 467
CLAESSON, M.H., RUDOLPHI, A., KOFOED, S., POULSEN, S.S. & REIMANN, J. $CD4^+$ T lymphocytes injected into severe combined immunodeficient (SCID) mice lead to an inflammatory and lethal bowel disease, 491
COCKWELL, P., ADAMS, D.H. & SAVAGE, C.O.S. Glycosaminoglycans contribute to multiple functions of vascular endothelial cells, 1
COLL-VINENT, B. *see* CID, M.-C.
COLOMBEL, J.F. *see* DESREUMAUX, P.
COLOMBEL, M. *see* LEVY, Y.
COMACCHIO, R.M. *see* WILLIAMS, W.M.
CORTOT, A. *see* DESREUMAUX, P.
CSERNOK, E. *see* STOFFEL, M.P.
CUADRADO, M.J. *see* GARCÍA-CÓZAR, F.J.
CUKROWSKA, B. *see* TUČKOVÁ, L.
CURRENTI, M. *see* ANTONELLI, G.
CZERKINSKY, C. *see* AHLFORS, E.

D'CRUZ, P.L. *see* MERONI, D.
D'ORAZIO, J.A. & STEIN-STREILEIN, J. Human natural killer (NK) cells present staphylococcal enterotoxin B (SEB) to T lymphocytes, 366
D'OSTILIO, N., SABATINO, G., NATOLI, C., ULLRICH, A. & IACOBELLI, S. 90K (Mac-2 BP) in human milk, 543
DANIEL, V., SÜSAL, C., WEIMER, R., ZIPPERLE, S., KRÖPELIN, M., ZIMMERMANN, R., HUTH-KÜHNE, A. & OPELZ, G. Association of T cell dysfunction with the presence of IgG autoantibodies on $CD4^+$ lymphocytes in haemophilia patients; results of a 10-year study, 4
DARK, J.H. *see* RIX, D.A.
DAVID, D. *see* DEMAISON, C.
DAVIES, K.A. *see* BOTTO, M.
DAVIES, K.A. *see* HOGARTH, M.B.
DE GEUS, B. *see* LUCA, M.E.
DE HEER, E. *see* LUCA, M.E.
DEL VECCHIO, S. *see* ANTONELLI, G.
DEMAISON, C., DAVID, D., FAUTREL, B. & THEZE, J. V_H gene-family

representation in peripheral activated B cells from systemic lupus erythematosus (SLE) patients, 439
DESREUMAUX, P., LEVY, A., VAERMAN, J.P., CAPRON, M., CORTOT, A. & COLOMBEL, J.F. Effect of intrajejunal elemental diet perfusion on local secretion of soluble CD4 and CD8, 293
DIANZANI, F. *see* ANTONELLI, G.
DIGHIERO, G. *see* CAYOTA, A.
DOE, W.F. *see* PAVLI, P.
DOHERTY, P.J. *see* MOLFINO, N.A.
DOMINGUEZ, M. *see* LIDINGTON, E.
DOMSCHKE, W. *see* KUCHARZIK, T.
DOUGLAS, M.S. *see* RIX, D.A.
DU PONT, J.S. *see* RICHTERS, C.D.

EGNER, W. *see* WONG, V.S.
EGUCHI, K. *see* MATSUOKA, N.
EGUCHI, K. *see* MIGITA, K.
ELSEY, T. *see* WONG, V.S.
ESAGUY, N. & AGUAS, A.P. Prevention of adjuvant arthritis in Lewis rats by neonatal bacille Calmette–Guérin (BCG) infection, 103
ESIN, S., BATONI, G., KÄLLENIUS, G., GAINES, H., CAMPA, M., SVENSON, S.B., ANDERSSON, R. & WIGZELL, H. Proliferation of distinct human T cell subsets in response to live, killed or soluble extracts of *Mycobacterium tuberculosis* and *Myco.avium*, 419
ESPARZA, J. *see* CID, M.-C.
EZAKI, I., OKADA, M., YOSHIKAWA, Y., FUJIKAWA, Y., HASHIMOTO, M., OTSUKA, M., NOMURA, T., YAMAMOTO, K., WATANABE, T., SHINGU, M. & NOBUNAGA, M. Human monoclonal rheumatoid factors augment arthritis in mice by the activation of T cells, 474

FAGERBERG, J. *see* WADHWA, M.
FAN, W.S. *see* LI, J.-M.
FAUTREL, B. *see* DEMAISON, C.
FAY, M. *see* RAJAGOPALAN-LEVASSEUR, P.
FELL, J.M.E., WALKER-SMITH, J.A., SPENCER, J. & MACDONALD, T.T. The distribution of dividing T cells throughout the intestinal wall in inflammatory bowel disease (IBD), 280
FERRY, B. *see* LIDINGTON, E.
FOURNIÉ, G.J. Detection of nucleosome–IgG immune complexes in ascites from mice transplanted with anti-DNA antibody-secreting hybridomas and in plasmas from MRL-*lpr*/*lpr* mice, 236
FRIMAN, V., HANSON, L.Å., BRIDON, J.-M., TARKOWSKI, A., BANCHEREAU, J. & BRIÈRE, F. IL-10-driven immunoglobulin production by B lymphocytes from IgA-deficient individuals correlates to infection proneness, 432
FUJIGAKI, Y. *see* MORIOKA, T.
FUJIKAWA, Y. *see* EZAKI, I.
FUJITA, T. *see* UEKI, T.

GAINES, H. *see* ESIN, S.
GAINES-DAS, R. *see* WADHWA, M.
GARCÍA-CÓZAR, F.J., MOLINA, I.J., CUADRADO, M.J., MARUBAYASHI, M., PEÑA, J. & SANTAMARÍA, M. Defective B7 expression on antigen-presenting cells underlying T cell activation abnormalities in systemic lupus erythematosus (SLE) patients, 72
GEJYO, F. *see* SUZUKI, S.
GHALIB, H.W. *see* SATTI, M.Z.
GIANNELLI, G. *see* ANTONELLI, G.
GLEICH, G.J. *see* OHNISHI, T.
GONZALEZ, G. *see* CAYOTA, A.
GORDON, T.P. *see* KEECH, C.L.
GOUGEROT-POCIDALO, M.-A. *see* RAJAGOPALAN-LEVASSEUR, P.
GRAU, J.-M. *see* CID, M.-C.
GREENSTEIN, J.D., PEAKE, P.W. & CHARLESWORTH, J.A. The metabolism of C9 in normal subjects and in patients with autoimmune disease, 160
GRINYÓ, J.M. *see* MESTRE, M.
GROSS, W.L. *see* STOFFEL, M.P.

HAAHEIM, L.R. *see* RØSOK, B.
HAMMOND, K.R. *see* KUTTEH, W.H.
HANIKÝŘOVÁ, M. *see* TUČKOVÁ, L.
HANSON, L.Å. *see* FRIMAN, V.
HASHIMOTO, M. *see* EZAKI, I.
HASHIMOTO, S. *see* NAKAYAMA, T.
HENTATI, B. *see* MATSIOTA-BERNARD, P.
HERNANDEZ-GARAY, M. *see* MÉNDEZ-SAMPERIO, P.
HERZBERG, C. *see* STOFFEL, M.P.
HOEKSTRA, M.J. *see* RICHTERS, C.D.
HOFFMAN, W.H. *see* JAUME, J.C.
HOFFMANN, J.C., BAYER, B. & ZEIDLER, H. Characterization of a soluble form of CD58 in synovial fluid of patients with rheumatoid arthritis (RA), 460
HOGARTH, M.B., NORSWORTHY, P.J., ALLEN, P.J., TRINDER, P.K.E., LOOS, M., MORLEY, B.J., WALPORT, M.J. & DAVIES, K.A. Autoantibodies to the collagenous region of C1q occur in three strains of lupus-prone mice, 241
HOLM, G. *see* YI, Q.
HOMEIDA, M.M.A. *see* SATTI, M.Z.
HORIE, T. *see* NAKAYAMA, T.
HORSFALL, A.C. *see* LI, J.-M.
HORST, G. *see* SPRONK, P.E.
HUBSCHER, O. *see* BACMAN, S.
HUGHES, G.R.V. *see* MERONI, D.
HUTH-KÜHNE, A. *see* DANIEL, V.
HYÖTY, H. *see* LÖNNROT, M.
HÖFLICH, C. *see* KERN, F.

IACOBELLI, S. *see* D'OSTILIO, N.
ILONEN, J. *see* MARTTILA, J.
INABA, T. *see* UEKI, T.
INMAN, R.D. & CHIU, B. Comparative microbicidal activity of synovial fluid on arthritogenic organisms, 80
ISENBERG, D.A. *see* BOTTO, M.
ISENBERG, D.A. *see* WILLIAMS, W.M.

JAUME, J.C., BUREK, C.L., HOFFMAN, W.H., ROSE, N.R., MCLACHLAN, S.M. & RAPOPORT, B. Thyroid peroxidase autoantibody epitopic 'fingerprints' in juvenile Hashimoto's thyroiditis: evidence for conservation over time and in families, 115
JOHNSTON, T. *see* STOFFEL, M.P.
JONSSON, R. *see* AHLFORS, E.
JOSÉ CAMUSSO, J. *see* BACMAN, S.
JOUVENNE, P. *see* MENETRIER-CAUX, C.

KAKIMOTO, K. *see* SAKATA, A.
KALLENBERG, C.G.M. *see* SPRONK, P.E.
KAMIYASU, M. *see* MYOZAKI, M.
KAMPERDIJK, E.W.A. *see* RICHTERS, C.D.
KANTELE, A. *see* KANTELE, J.M.
KANTELE, J.M., KANTELE, A. & ARVILOMMI, H. Circulating immunoglobulin-secreting cells are heterogeneous in their expression of maturation markers and homing receptors, 525
KAWABE, Y. *see* MATSUOKA, N.
KAWABE, Y. *see* MIGITA, K.
KAWAKAMI, A. *see* MATSUOKA, N.
KAWAKAMI, K., TOHYAMA, M., XIE, Q. & SAITO, A. IL-12 protects mice against pulmonary and disseminated infection caused by *Cryptococcus neoformans*, 208
KEECH, C.L., GORDON, T.P. & MCCLUSKEY, J. Structural differences between the human and mouse 52-kD Ro autoantigens associated with poorly conserved autoantibody activity across species, 255
KEKWICK, R., BHAMBRI, S., CHABANE, M.H., AUTEGARDEN, J.-E., LEVY, D.A. & LEYNADIER, F. The allergenic properties of fresh and preserved *Hevea brasiliensis* latex protein preparations, 337
KERN, F., ODE-HAKIM, S., VOGT, K., HÖFLICH, C., REINKE, P. & VOLK, H.D. The enigma of $CD57^+CD28^-$ T cell expansion—anergy or activation?, 180
KESTEN, S. *see* MOLFINO, N.A.
KHAMASHTA, M. *see* MERONI, D.
KIHARA, Y. *see* UEKI, T.
KIRBY, J.A. *see* RIX, D.A.
KISO, T. *see* UEKI, T.
KITA, H. *see* OHNISHI, T.
KNIP, M. *see* LÖNNROT, M.
KOFOED, S. *see* CLAESSON, M.H.
KOMATSU, T., MORIYA, N. & SHIOHARA, T. T cell receptor (TCR) repertoire and function of human epidermal T cells: restricted TCR

Vα-Vβ genes are utilized by T cells residing in the lesional epidermis in fixed drug eruption, 343
KRAAL, G. *see* BROUG-HOLUB, E.
KRÖPELIN, M. *see* DANIEL, V.
KUCHARZIK, T., LÜGERING, N., WEIGELT, H., ADOLF, M., DOMSCHKE, W. & STOLL, R. Immunoregulatory properties of IL-13 in patients with inflammatory bowel disease; comparison with IL-4 and IL-10, 483
KULMALA, P. *see* LÖNNROT, M.
KUTTEH, C.C. *see* KUTTEH, W.H.
KUTTEH, W.H., PRINCE, S.J. HAMMOND, K.R., KUTTEH, C.C. & MESTECKY, J. : Variations in immunoglobulins and IgA subclasses of human uterine cervical secretions around the time of ovulation, 538
KÄLLENIUS, G. *see* ESIN, S.

LABAUME, S. *see* LEVY, Y.
LARSSON, E. *see* LI, J.-M.
LATOUF, S.E. *see* LUKEY, P.T.
LECOINTE, D. *see* RAJAGOPALAN-LEVASSEUR, P.
LEFVERT, A.K. *see* YI, Q.
LEGAKIS, N. *see* MATSIOTA-BERNARD, P.
LEHTINEN, M. *see* MARTTILA, J.
LEINIKKI, P. *see* LÖNNROT, M.
LEIRISALO-REPO, M. *see* VAARALA, O.
LEONI, J. *see* ABATANGELO, C.
LEVY, A. *see* DESREUMAUX, P.
LEVY, D.A. *see* KEKWICK, R.
LEVY, Y., LABAUME, S., COLOMBEL, M. & BROUET, J.C. Retinoic acid modulates the *in vivo* and *in vitro* growth of IL-6 autocrine human myeloma cell lines via induction of apoptosis, 167
LI, J.-M., FAN, W.S., HORSFALL, A.C., ANDERSON, A.C., RIGBY, S., LARSSON, E. & VENABLES, P.J.W. The expression of human endogenous retrovirus-3 in fetal cardiac tissue and antibodies in congenital heart block, 388
LIDINGTON, E., NÖHAMMER, C., DOMINGUEZ, M., FERRY, B. & ROSE, M.L. Inhibition of the transendothelial migration of human lymphocytes but not monocytes by phosphodiesterase inhibitors, 66
LIMBURG, P.C. *see* SPRONK, P.E.
LINK, H. *see* MA, C.-G.
LOOS, M. *see* HOGARTH, M.B.
LÖNNROT, M., HYÖTY, H., KNIP, M., ROIVAINEN, M., KULMALA, P., LEINIKKI, P., ÅKERBLOM, H.K. and the Childhood Diabetes in Finland Study Group: Antibody cross-reactivity induced by the homologous regions in glutamic acid decarboxylase (GAD_{65}) and 2C protein of coxsackievirus B4, 398
LUCA, M.E., DE GEUS, B., SAHALI, D., BRUIJN, J.A., VERROUST, P. & DE HEER, E. Isolation of cDNAs encoding immunogenic regions of gp330, the autoantigen involved in Heymann nephritis, 312
LÜGERING, N. *see* KUCHARZIK, T.
LUKEY, P.T., LATOUF, S.E. & RESS, S.R. Memory lymphocytes from tuberculous effusions: purified protein derivative (PPD) stimulates accelerated activation marker expression and cell cycle progression, 412
LUKKA, M. *see* VAARALA, O.

MA, C.-G., ZHANG, G.-X., XIAO, B.-G. & LINK, H. Cellular mRNA expression of interferon-gamma (IFN-γ), IL-4 and transforming growth factor-beta (TGF-β) in rats nasally tolerized against experimental autoimmune myasthenia gravis (EAMG), 509
MACDONALD, T.T. *see* FELL, J.M.E.
MAGGI, F. *see* ANTONELLI, G.
MARTTILA, J., ILONEN, J., LEHTINEN, M., PARKKONEN, P. & SALMI, A. Definition of three minimal T helper cell epitopes of rubella virus E1 glycoprotein, 394
MARUBAYASHI, M. *see* GARCÍA-CÓZAR, F.J.
MARUYAMA, N. *see* TAKAHASHI, T.
MASCAGNI, B. *see* TEDESCHI, A.
MASSIP, E. *see* MESTRE, M.
MATHOV, I. *see* ABATANGELO, C.
MATSIOTA-BERNARD, P., HENTATI, B., PIÉ, S., LEGAKIS, N., NAUCIEL, C. & AVRAMEAS, S. Beneficial effect of *Salmonella typhimurium* infection and of immunoglobulins from *S.typhimurium*-infected mice on the autoimmune disease of (NZB × NZW)F_1 mice, 228
MATSUMURA, O. *see* TAKAHASHI, T.
MATSUOKA, N., EGUCHI, K., KAWAKAMI, A., TSUBOI, M., KAWABE, Y., AOYAGI, T. & NAGATAKI, S. Inhibitory effect of clarithromycin on costimulatory molecule expression and cytokine production by synovial fibroblast-like cells, 501
MATSUOKA, Y. *see* UEKI, T.
MAXWELL, L. *see* PAVLI, P.
MAYENO, A.N. *see* OHNISHI, T.
MC CUTCHAN, J.A. *see* WILLIAMS, W.M.
MCCLUSKEY, J. *see* KEECH, C.L.
MCKILLOP, J.H. *see* CHEN, G.
MCLACHLAN, S.M. *see* JAUME, J.C.
MEDINA, E. & NORTH, R.J. Mice that carry the resistance allele of the *Bcg* gene (*Bcg*^{r}) develop a superior capacity to stabilize bacille Calmette–Guérin (BCG) infection in their lungs and spleen over a protracted period in the absence of specific immunity, 44
MELLSTEDT, H. *see* WADHWA, M.
MÉNDEZ-SAMPERIO, P., HERNANDEZ-GARAY, M., BADILLO-FLORES, A. & NUÑEZ-VAZQUEZ, A. Down-modulation of mycobacterial-induced IL-1β production in human mononuclear cells by IL-4, 374
MENDIS, K.N. *see* WIJESEKERA, S.K.
MENETRIER-CAUX, C., BRIERE, F., JOUVENNE, P., PEYRON, E., PEYRON, F. & BANCHEREAU, J. Identification of human IgG autoantibodies specific for IL-10, 173
MERONI, D. D'CRUZ, P.L., KHAMASHTA, M., YOUINOU, P. & HUGHES, G.R.V. Anti-endothelial cell antibodies: only for scientists or for clinicians too?, 199
MESTECKY, J. *see* KUTTEH, W.H.
MESTRE, M., MASSIP, E., BAS, J., ALSINA, J., ROMEU, A., CASTELAO, A.M., BUENDIA, E. & GRINYÓ, J.M. Longitudinal study of the frequency of cytotoxic T cell precursors in kidney allograft recipients, 108
MEZHLUMOVA, M.B. *see* NIKONENKO, B.V.
MIADONNA, A. *see* TEDESCHI, A.
MIGITA, K., EGUCHI, K., AOYAGI, T., TSUKADA, T., TSUBOI, M., KAWABE, Y. & NAGATAKI, S. The effects of the immunosuppressant rapamycin on the growth of rheumatoid arthritis (RA) synovial fibroblast , 86
MILAZZO, N. *see* TEDESCHI, A.
MISRA, A., RUMA MUKERJEE & CHATURVEDI, U.C. Production of nitrite by dengue virus-induced cytotoxic factor, 406
MITARAI, T. *see* TAKAHASHI, T.
MIURA, T. *see* MYOZAKI, M.
MIYAKE, K. *see* TAKAHASHI, T.
MIZUNO, M. *see* UEKI, T.
MOLFINO, N.A., DOHERTY, P.J., SUURMANN, I.L., YANG, S.X., KESTEN, S., CHAPMAN, K.R. & SLUTSKY, A.S. Analysis of the T cell receptor Vγ region gene repertoire in bronchoalveolar lavage (BAL) and peripheral blood of atopic asthmatics and healthy subjects, 144
MOLINA, I.J. *see* GARCÍA-CÓZAR, F.J.
MORGAN, L. *see* ZHANG, J.-G.
MORIOKA, T., FUJIGAKI, Y., BATSFORD, S.R., WOITAS, R., OITE, T., SHIMIZU, F. & VOGT, A. Anti-DNA antibody derived from a systemic lupus erythematosus (SLE) patient forms histone–DNA–anti-DNA complexes that bind to rat glomeruli *in vivo*, 92
MORIYA, N. *see* KOMATSU, T.
MORLEY, B.J. *see* HOGARTH, M.B.
MOROZ, A.M. *see* NIKONENKO, B.V.
MORROW, W.J.W. *see* WILLIAMS, W.M.
MYOZAKI, M., KAMIYASU, M., MIURA, T., WATANABE, Y., NAKANISHI, T. & YAMASHITA, U. Induction of autoimmune hepatitis and autoantibodies to liver antigens by neonatal thymectomy in mice, 133

NAGATAKI, S. *see* MATSUOKA, N.
NAGATAKI, S. *see* MIGITA, K.
NAKANISHI, T. *see* MYOZAKI, M.
NAKATOMI, Y. *see* SUZUKI, S.
NAKAYAMA, T. HASHIMOTO, S. AMEMIYA, E. & HORIE, T. Elevation of plasma-soluble tumour necrosis factor receptors (TNF-R) in sarcoidosis, 318
NASU, J. *see* UEKI, T.
NATOLI, C. *see* D'OSTILIO, N.
NATORI, Y. *see* TAKAHASHI, T.

NAUCIEL, C. *see* MATSIOTA-BERNARD, P.
NIKONENKO, B.V., APT, A.S., MEZHLUMOVA, M.B., AVDIENKO, V.G., YEREMEEV, V.V. & MOROZ, A.M. Influence of the mouse *Bcg*, *Tbc-1* and *xid* genes on resistance and immune responses to tuberculosis infection and efficacy of bacille Calmette–Guérin (BCG) vaccination, 37
NOBUNAGA, M. *see* EZAKI, I.
NÖHAMMER, C. *see* LIDINGTON, E.
NOMURA, T. *see* EZAKI, I.
NORSWORTHY, P.J. *see* HOGARTH, M.B.
NORTH, R.J. *see* MEDINA, E.
NUÑEZ-VAZQUEZ, A. *see* MÉNDEZ-SAMPERIO, P.

ODANI, S. *see* SUZUKI, S.
ODE-HAKIM, S. *see* KERN, F.
OHMURA, T. *see* SAKATA, A.
OHNISHI, T., KITA, H., MAYENO, A.N., OKADA, S., SUR, S., BROIDE, D.H. & GLEICH, G.J. Lidocaine in bronchoalveolar lavage fluid (BALF) is an inhibitor of eosinophil-active cytokines, 325
OITE, T. *see* MORIOKA, T.
OKADA, H. *see* UEKI, T.
OKADA, M. *see* EZAKI, I.
OKADA, S. *see* OHNISHI, T.
OPELZ, G. *see* DANIEL, V.
OTSUKA, M. *see* EZAKI, I.
OTTENHOFF, T.H.M. Putting numbers on mycobacterium activated T cell subsets, 381

PALELLA, M. *see* TEDESCHI, A.
PALOSUO, T. *see* VAARALA, O.
PARKKONEN, P. *see* MARTTILA, J.
PASTORE, G. *see* ANTONELLI, G.
PAVLI, P., MAXWELL, L., VAN DE POL, E. & DOE, W.F. Distribution of human colonic dendritic cells and macrophages, 124
PEAKE, P.W. *see* GREENSTEIN, J.D.
PEDROL, E. *see* CID, M.-C.
PEÑA, J. *see* GARCÍA-CÓZAR, F.J.
PEYRON, E. *see* MENETRIER-CAUX, C.
PEYRON, F. *see* MENETRIER-CAUX, C.
PICAZO, A. *see* CID, M.-C.
PIÉ, S. *see* MATSIOTA-BERNARD, P.
PING, H. *see* SAKATA, A.
PISTELLO, M. *see* ANTONELLI, G.
PLOTKIN, L. *see* ABATANGELO, C.
POULSEN, S.S. *see* CLAESSON, M.H.
PRINCE, S.J. *see* KUTTEH, W.H.
PUURUNEN, M. *see* VAARALA, O.
PYRAH, I.T.G. & WATT, N.J. Immunohistological study of the depressed cutaneous DTH response in sheep naturally infected with an ovine lentivirus (Maedi-Visna virus), 32

RAGNHAMMAR, P. *see* WADHWA, M.
RAJAGOPALAN-LEVASSEUR, P., LECOINTE, D., BERTRAND, G., FAY, M. & GOUGEROT-POCIDALO, M.-A. Differential nitric oxide (NO) production by macrophages from mice and guinea pigs infected with virulent and avirulent *Legionella pneumophila* serogroup 1, 48
RAPOPORT, B. *see* JAUME, J.C.
RATHNAYAKA, L. *see* WIJESEKERA, S.K.
REIMANN, J. *see* CLAESSON, M.H.
REINKE, P. *see* KERN, F.
REITS, E.A.J. *see* RICHTERS, C.D.
RESS, S.R. *see* LUKEY, P.T.
RICHTERS, C.D., REITS, E.A.J., VAN PELT, A.M., HOEKSTRA, M.J., VAN BAARE, J., DU PONT, J.S. & KAMPERDIJK, E.W.A. Effect of low dose UVB irradiation on the migratory properties and functional capacities of human skin dendritic cells, 191
RIGBY, S. *see* LI, J.-M.
RIX, D.A., DOUGLAS, M.S., TALBOT, D., DARK, J.H. & KIRBY, J.A. Role of glycosaminoglycans (GAGs) in regulation of the immunogenicity of human vascular endothelial cells, 60
ROFFI, L. *see* ANTONELLI, G.
ROIVAINEN, M. *see* LÖNNROT, M.
ROMEU, A. *see* MESTRE, M.
ROSE, M.L. *see* LIDINGTON, E.
ROSE, N.R. *see* JAUME, J.C.
ROSENBERG, J. *see* WILLIAMS, W.M.
RØSOK, B. VOLTERSVIK, P. BJERKNES, R. AXELSSON, M. HAAHEIM, L.R. & ÅSJÖ, B. Dynamics of HIV-1 replication following influenza vaccination of HIV^+ individuals, 203
ROSS, G.D. *see* THORNTON, B.P.
RUDOLPHI, A. *see* CLAESSON, M.H.

SABATINO, G. *see* D'OSTILIO, N.
SAHALI, D. *see* LUCA, M.E.
SAITO, A. *see* KAWAKAMI, K.
SAKAI, O. *see* TAKAHASHI, T.
SAKATA, A., SAKATA, K., PING, H., OHMURA, T., TSUKANO, M. & KAKIMOTO, K. Successful induction of severe destructive arthritis by the transfer of *in vitro*-activated synovial fluid T cells from patients with rheumatoid arthritis (RA) in severe combined immunodeficient (SCID) mice, 247
SAKATA, K. *see* SAKATA, A.
SALMI, A. *see* MARTTILA, J.
SANTAMARÍA, M. *see* GARCÍA-CÓZAR, F.J.
SATO, H. *see* SUZUKI, S.
SATTI, M.Z., SULAIMAN, S.M., HOMEIDA, M.M.A., YOUNIS, S.A. & GHALIB, H.W. Clinical, parasitological and immunological features of canal cleaners hyper-exposed to *Schistosoma mansoni* in the Sudan, 426
SAVAGE, C.O.S. *see* COCKWELL, P.
SHIMIZU, F. *see* MORIOKA, T.
SHINGU, M. *see* EZAKI, I.
SHIOHARA, T. *see* KOMATSU, T.
SHIRASAWA, T. *see* TAKAHASHI, T.
SIMEONI, E. *see* ANTONELLI, G.
SLUTSKY, A.S. *see* MOLFINO, N.A.
SPENCER, J. *see* FELL, J.M.E.
SPICKETT, G.P. *see* ZHANG, J.-G.
SPRONK, P.E., HORST, G., VAN DER GUN, B.T.F., LIMBURG, P.C. & KALLENBERG, C.G.M. Anti-dsDNA production coincides with concurrent B and T cell activation during development of active disease in systemic lupus erythematosus (SLE), 446
SQUIQUERA, L. *see* ABATANGELO, C.
STEIN-STREILEIN, J. *see* D'ORAZIO, J.A.
STERIN-BORDA, L. *see* BACMAN, S.
STOFFEL, M.P., CSERNOK, E., HERZBERG, C., JOHNSTON, T., CARROLL, S.F. & GROSS, W.L. Anti-neutrophil cytoplasmic antibodies (ANCA) directed against bactericidal/permeability increasing protein (BPI): a new seromarker for inflammatory bowel disease and associated disorders, 54
STOLL, R. *see* KUCHARZIK, T.
SULAIMAN, S.M. *see* SATTI, M.Z.
SUR, S. *see* OHNISHI, T.
SUURMANN, I.L. *see* MOLFINO, N.A.
SUZUKI, S., NAKATOMI, Y., ODANI, S., SATO, H., GEJYO, F. & ARAKAWA, M. Circulating IgA, IgG, and IgM class antibody against *Haemophilus parainfluenzae* antigens in patients with IgA nephropathy, 306
SVENSON, S.B. *see* ESIN, S.
SÜSAL, C. *see* DANIEL, V.

TAKAHASHI, T., SHIRASAWA, T., MIYAKE, K., YAHAGI, Y., MATSUMURA, O., ABE, A., NATORI, Y., YAMABE, H., UTSUNOMIYA, Y., MARUYAMA, N., MITARAI, T. & SAKAI, O. Jak3 expression in glomerular epithelia of IgA nephropathy (IgA-N) patients, 517
TALBOT, D. *see* RIX, D.A.
TARKOWSKI, A. *see* FRIMAN, V.
TEDESCHI, A., PALELLA, M., MILAZZO, N., MASCAGNI, B. & MIADONNA, A. Impairment of the inhibitory effect of sodium on basophil histamine release in patients with systemic sclerosis, 97
THEODORIDIS, E. *see* BOTTO, M.
THEZE, J. *see* DEMAISON, C.
THOMPSON, E.M. *see* BOTTO, M.
THORNTON, B.P., VĚTVIČKA, V. & ROSS, G.D. Function of C3 in a humoral response: iC3b/C3dg bound to an immune complex generated with natural antibody and a primary antigen promotes antigen uptake and the expression of co-stimulatory molecules by all B cells, but only stimulates immunoglobulin synthesis by antigen-specific B cells, 531

THORPE, R. *see* WADHWA, M.
TLASKALOVÁ-HOGENOVÁ, H. *see* TUČKOVÁ, L.
TOBÍAS, E. *see* CID, M.-C.
TOHYAMA, M. *see* KAWAKAMI, K.
TRINDER, P.K.E. *see* HOGARTH, M.B.
TSUBOI, M. *see* MATSUOKA, N.
TSUBOI, M. *see* MIGITA, K.
TSUJI, T. *see* UEKI, T.
TSUKADA, T. *see* MIGITA, K.
TSUKANO, M. *see* SAKATA, A.
TUČKOVÁ, L., ZÍDEK, Z., HANIKÝŘOVÁ, M., CUKROWSKA, B., TLASKALOVÁ-HOGENOVÁ, H. & BAROT-CIORBARU, R. Macrophage nitric oxide synthase (NOS) activation by *Nocardia opaca* fractions and 15- and 56-kD isolated antigens, 215

UEKI, T., MIZUNO, M., UESU, T., KISO, T., NASU, J., INABA, T., KIHARA, Y., MATSUOKA, Y., OKADA, H., FUJITA, T. & TSUJI, T. Distribution of activated complement, C3b, and its degraded fragments, iC3b/C3dg, in the colonic mucosa of ulcerative colitis (UC), 286
UESU, T. *see* UEKI, T.
ULLRICH, A. *see* D'OSTILIO, N.
URBANO-MÁRQUEZ, A. *see* CID, M.-C.
UTSUNOMIYA, Y. *see* TAKAHASHI, T.

VAARALA, O., PUURUNEN, M., LUKKA, M., ALFTHAN, G., LEIRISALO-REPO, M., AHO, K. & PALOSUO, T. Affinity-purified cardiolipin-binding antibodies show heterogeneity in their binding to oxidized low-density lipoprotein, 269
VAERMAN, J.P. *see* DESREUMAUX, P.
VAN BAARE, J. *see* RICHTERS, C.D.
VAN DE POL, E. *see* PAVLI, P.
VAN DER GUN, B.T.F. *see* SPRONK, P.E.
VAN PELT, A.M. *see* RICHTERS, C.D.
VENABLES, P.J.W. *see* LI, J.-M.
VERROUST, P. *see* LUCA, M.E.
VĚTVIČKA, V. *see* THORNTON, B.P.
VOGT, A. *see* MORIOKA, T.
VOGT, K. *see* KERN, F.
VOLK, H.D. *see* KERN, F.
VOLTERSVIK, P. *see* RØSOK, B.
VUILLIER, F. *see* CAYOTA, A.

WADHWA, M., BIRD, C., FAGERBERG, J., GAINES-DAS, R., RAGNHAMMAR, P., MELLSTEDT, H. & THORPE, R. Production of neutralizing granulocyte-macrophage colony-stimulating factor (GM-CSF) antibodies in carcinoma patients following GM-CSF combination therapy, 351
WALKER, J.J. *see* CHEN, G.
WALKER-SMITH, J.A. *see* FELL, J.M.E.
WALPORT, M.J. *see* BOTTO, M.
WALPORT, M.J. *see* HOGARTH, M.B.
WANG, S.H. *see* CHEN, G.
WATANABE, T. *see* EZAKI, I.
WATANABE, Y. *see* MYOZAKI, M.
WATT, N.J. *see* PYRAH, I.T.G.
WATTS, R.A. *see* WILLIAMS, W.M.
WEIGELT, H. *see* KUCHARZIK, T.
WEIMER, R. *see* DANIEL, V.
WHALLEY, A.S. *see* WILLIAMS, W.M.
WIGZELL, H. *see* ESIN, S.
WIJESEKERA, S.K., CARTER, R., RATHNAYAKA, L. & MENDIS, K.N. A malaria parasite toxin associated with *Plasmodium vivax* paroxysms, 221
WILLIAMS, W.M., WHALLEY, A.S., COMACCHIO, R.M. ROSENBERG, J., WATTS, R.A., ISENBERG, D.A., MCCUTCHAN, J.A. & MORROW, W.J.W. : Correlation between expression of antibodies to histone H2B and clinical activity in HIV-infected individuals, 18
WILSON, R. *see* CHEN, G.
WOITAS, R. *see* MORIOKA, T.
WONG, V.S. EGNER, W., ELSEY, T., BROWN, D. & ALEXANDER, G.J.M. Incidence, character and clinical relevance of mixed cryoglobulinaemia in patients with chronic hepatitis C virus infection, 25

XIAO, B.-G. *see* MA, C.-G.
XIE, Q. *see* KAWAKAMI, K.

YAHAGI, Y. *see* TAKAHASHI, T.
YAMABE, H. *see* TAKAHASHI, T.
YAMAMOTO, K. *see* EZAKI, I.
YAMASHITA, U. *see* MYOZAKI, M.
YANG, S.X. *see* MOLFINO, N.A.
YEREMEEV, V.V. *see* NIKONENKO, B.V.
YI, Q., HOLM, G. & LEFVERT, A.K. Idiotype-induced T cell stimulation requires antigen presentation in association with HLA-DR molecules, 359
YOSHIKAWA, Y. *see* EZAKI, I.
YOUINOU, P. *see* MERONI, D.
YOUNIS, S.A. *see* SATTI, M.Z.

ZEIDLER, H. *see* HOFFMANN, J.C.
ZHANG, G.-X. *see* MA, C.-G.
ZHANG, J.-G., MORGAN, L. & SPICKETT, G.P. L-selectin in patients with common variable immunodeficiency (CVID): a comparative study with normal individuals, 275
ZHENG, H.Z. *see* CHEN, G.
ZÍDEK, Z. *see* TUČKOVÁ, L.
ZIMMERMANN, R. *see* DANIEL, V.
ZIPPERLE, S. *see* DANIEL, V.

Subject Index

52-kD Ro, 255

adhesion, 66
adhesion molecule, 501
adhesion molecules, 467
alveolar macrophages, 332
anti-C1q antibodies, 241
anti-DNA, 439
anti-DNA antibody, 92
anti-dsDNA, 446
anti-lymphocyte antibodies, 18
anti-lymphocyte autoantibodies, 4
anti-neutrophil cytoplasmic antibodies (ANCA), 54
antibodies, 103, 454
antibodies to interferon, 384
antibody, 236
antibody production, 351
anticardiolipin antibodies, 269
antigen presentation, 359
antiphospholipid antibodies, 269
apoptosis, 167
arthritis model, 247
asthma, 144
autoantibodies, 173
autoantibody, 115, 133
autoantigens, 255
autoimmune disease, 160
autoimmune diseases, 388
autoimmune hepatitis, 133
autoimmunity, 18, 103, 185, 439

β2-glycoprotein I, 269
B cells, 275
B lymphocytes, 432, 531
B7 molecules, 191
B7/CD28, 72
bacteria, 80
bactericidal/permeability increasing protein, 54
basophil, 97
Bcg gene, 44
BCG infection, 44
bronchial asthma, 325
bronchoalveolar lavage fluid, 325
BXSB, 241

C3, 531
C3b, 286
C9, 160
canal cleaners, 426
CD2, 460
CD28 antigen, 180
$CD4^+$ lymphocytes, 4, 11
CD40, 432
CD45, 11
CD57 antigen, 180
CD58, 460
CD80, 531
cDNA, 312
CDR1H, 185
cell cycle, 412
cell death, 236
cercarial homogenate, 426
cervical mucus, 538
clarithromycin, 501
class II MHC antigen, 60
class switch, 306
clinical effects, 351
clinical immunity, 221
clonal deletion, 108
cold agglutinins, 185
collagen-induced arthritis, 474
common variable immunodeficiency, 175
complement, 25, 286
coxsackievirus B4, 398
CR2/CD21, 531
Crohn's disease, 280
cross-reactivity, 351, 398
cytokine, 501
cytokines, 332
cytomegalovirus infection, 180
cytotoxic factor, 406
cytotoxicity, 406

delayed-type hypersensitivity, 297
dendritic cells, 124
dengue virus, 406
dissemination, 208
drug eruption, 343

E1 glycoprotein, 394
effect of immunoglobulins, 228
elemental diet, 293
ELISPOT, 525
endothelium, 60, 66
eosinophil survival activity, 325
epidermal T cells, 343
epitope, 115
experimental autoimmune myasthenia gravis, 509

Fc receptor, 264
fingerprint, 115

γ_c chain, 517
gastrointestinal infection, 543
gene expression, 154
gene polymorphism, 154
glomerular epithelium, 517
glucocorticoids, 332
glutamic acid decarboxylase, 398
glycosaminoglycan, 60
GM-CSF, 351
gp330, 312

haemagglutination inhibition, 203
haemophilia, 4
Haemophilus parainfluenzae, 306
HCV-RNA, 25
heat shock protein, 374
hepatitis C, 384
hepatitis C treatment, 384
hepatitis C virus, 25
Heymann nephritis, 312
histamine release, 97
histone, 92
histones, 18
HIV, 4, 11, 18, 203
homing receptors, 525
host defence, 543
hsp, 144
human, 160, 173

hypersensitivity, 32
hypertension, 154

iC3b/C3dg, 286
idiotype, 359
IFN-γ, 208, 509
IgA, 538
IgA deficiency, 432
IgA nephropathy, 306, 517
IL-1, 374
IL-10, 173, 432, 483
IL-12, 208
IL-13, 483
IL-4, 374, 483, 509
IL-5, 325
IL-6, 167
IL-6 receptor, 167
immune activation, 180
immune response, 37
immunoglobulin production, 432
immunoglobulin-secreting cells, 525
immunoglobulins, 538
immunohistochemistry, 297, 517
immunology, 32
immunomagnetic cell sorting, 525
inflammation, 144, 467
inflammatory bowel disease, 54, 483, 491
influenza, 203
insulin-dependent diabetes mellitus, 398
intercellular adhesion molecule-1, 60
interferon-alpha, 384
interferon-gamma, 60
interferons, 384
intestinal antigen handling, 124
intestinal macrophages, 124
intestinal perfusion, 293

Jak3, 517

kidney, 108

L-selectin, 275
lamina propria, 280
Langerhans cells, 191
latex allergens, 337
latex maturation, 337
Legionella pneumophila, 48
LFA-1/CD11a, 531
LFA-3, 191, 460
lidocaine, 325
limiting dilution assay, 108
liver antigens, 133
lung, 144
lupus, 236, 241, 439
lupus nephritis, 92
lymphocyte activation, 446
lymphocyte adhesion, 60
lymphocyte homing, 525

M-component, 359
M_3 muscarinic receptors, 454
macrophage activation, 215
macrophages, 48
Maedi-Visna, 32
malaria toxin, 221
maternal autoantibodies, 388
memory cells, 412
metabolism, 160
MHC molecules, 359
mice, 48
migration, 66, 191
mixed cryoglobulinaemia, 25
monoclonal anti-DNA, 236
monocytes, 483
mouse, 297
MRL/lpr, 241
mucosal immunity, 538
multiple myeloma, 167
muscle, 280
mycobacteria, 103, 374
Mycobacterium avium, 419
Mycobacterium bovis-BCG, 103
Mycobacterium tuberculosis, 419
myopathies, 467

natural killer cells, 366, 419
natural resistance, 44
neonatal thymectomy, 133
neutralization, 351
newborn, 543
nitric oxide, 48
nitric oxide synthase, 215
nitrite, 406
Nocardia opaca, 215
non-H-2 genes, 37
nucleosome, 92
nucleosome–IgG complexes, 236
nucleosomes, 236
(NZB × NZW)F_1 lupus prone mice, 228
NZB/W, 241

oral mucosa, 297
outer membrane, 306
oxidative stress, 11
oxidized low-density lipoprotein, 269

parotid gland, 454
paroxysms, 221
pathogenesis, 221
phosphodiesterase, 66
Plasmodium vivax malaria, 221
polymerase chain reaction, 343
polymorphism, 264
pregnancy, 154, 388
primary Sjögren's syndrome, 454
proliferation, 280, 419
prospective study, 446
protein kinase, 86
public idiotope, 185
pulmonary cryptococcosis, 208

rapamycin, 86
reactive arthritis, 80
resistance to interferon therapy, 384
retinoic acid, 167
rheumatoid arthritis, 86, 103, 247, 460, 474
rheumatoid factor, 474
Ro/SS-A, 255
rubella virus, 394

Salmonella typhimurium infection, 228
sarcoidosis, 318
sCD4, 293
sCD8, 293
Schistosoma mansoni, 426
SCID mice, 247
SCID mouse, 491
serosa, 280
sheep, 32
skin prick tests, 337
sodium, 97
soluble adhesion molecules, 460
soluble TNF receptor, 318
staphylococcal enterotoxin B, 366
superantigen, 439
synovial fibroblast, 86
synovial fluid T cells, 247
synoviocyte, 501

synthetic peptides, 185, 398
systemic lupus erythematosus, 72, 264, 269, 446
systemic sclerosis, 97

T cell dysfunction, 4
T cell, 280
T cell epitopes, 394
T cell-depleted mice, 44
T cells, 72, 144, 275, 359, 491
T lymphocytes, 366, 419
TCR Vα-Vβ genes, 343
thyroid peroxidase, 115
thyroiditis, 115
tolerance, 509
transforming growth factor-beta, 509
transplantation, 108
tuberculosis, 37, 374, 412
tumour necrosis factor, 154
type II collagen, 474
tyrosine phosphatase, 11

ulcerative colitis, 286

V gene family, 439
vaccination, 203
viral load, 203
virulence, 48

whole worn homogenate, 426